

Generating Functions

Prepared by Mark on June 5, 2026
Based on a handout by Aaron Anderson

Part 1: Introduction

Definition 1:

Say we have a sequence a_0, a_1, a_2, \dots

The *generating function* of this sequence is defined as follows:

$$A(x) = \sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \dots$$

Under some circumstances, this sum does not converge, and thus $A(x)$ is undefined.

However, we can still manipulate this infinite sum to get useful results even if $A(x)$ diverges.

Problem 2:

Let $A(x)$ be the generating function of the sequence a_n ,

and let $B(x)$ be the generating function of the sequence b_n .

Find the sequences that correspond to the following generating functions:

- $cA(x)$
- $xA(x)$
- $A(x) + B(x)$
- $A(x)B(x)$

Problem 3:

Assuming $|x| < 1$, show that

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots$$

Hint: use some clever algebra. What is $x \times (1 + x + x^2 + \dots)$?

Problem 4:

Let $A(x)$ be the generating function of the sequence a_n .

Find the sequence that corresponds to the generating function $\frac{A(x)}{1-x}$

Problem 5:

Find short expressions for the generating functions for the following sequences:

- 1, 0, 1, 0, ...
- 1, 2, 4, 8, 16, ...
- 1, 2, 3, 4, 5, ...

Part 2: Fibonacci

Definition 6:

The *Fibonacci numbers* are defined by the following recurrence relation:

- $f_0 = 0$
- $f_1 = 1$
- $f_n = f_{n-1} + f_{n-2}$

Problem 7:

Let $F(x)$ be the generating function that corresponds to the Fibonacci numbers.

Find the generating function of $0, f_0, f_1, \dots$ in terms of $F(x)$.

Call this $G(x)$.

Problem 8:

Find the generating function of $0, 0, f_0, f_1, \dots$ in terms of $F(x)$. Call this $H(x)$.

Problem 9:

Calculate $F(x) - G(x) - H(x)$ using the recurrence relation that we used to define the Fibonacci numbers.

Problem 10:

Using the problems on the previous page, find $F(x)$ in terms of x .

Definition 11:

A *rational function* f is a function that can be written as a quotient of polynomials. That is, $f(x) = \frac{p(x)}{q(x)}$ where p and q are polynomials.

Problem 12:

Solve the equation from ?? for $F(x)$, expressing it as a rational function.

Definition 13:

Partial fraction decomposition is an algebraic technique that works as follows:

If $p(x)$ is a polynomial of degree 1 and a and b are constants, we can rewrite the rational function $\frac{p(x)}{(x-a)(x-b)}$ as follows:

$$\frac{p(x)}{(x-a)(x-b)} = \frac{c}{x-a} + \frac{d}{x-b}$$

where c and d are constants.

Problem 14:

Now that we have a rational function for $F(x)$,

find a closed-form expression for its coefficients using partial fraction decomposition.

Problem 15:

Using problems from the introduction and ??, find an expression for the coefficients of $F(x)$ (and thus, for the Fibonacci numbers).

Problem 16: Bonus

Repeat the method of recurrence, generating function, partial fraction decomposition, and geometric series to find a closed form for the following sequence:

$$a_0 = 1; \quad a_{n+1} = 2a_n + n$$

Hint: When doing partial fraction decomposition with a denominator of the form $(x - a)^2(x - b)$, you may need to express your expression as a sum of three fractions: $\frac{c}{(x-a)^2} + \frac{d}{x-a} + \frac{e}{x-b}$.

Part 3: Dice

Definition 17:

A *die* is a device that randomly selects a positive integer from a finite list of options. For example, the standard 6-sided die selects a value from $[1, 2, 3, 4, 5, 6]$. We may have many sides with the same value, as in $[1, 1, 2, 3]$.

To describe a die with a generating function, let a_k be the number of times k appears as a side of the die and consider $a_0 + a_1x + a_2x^2 + \dots$

A die has a finite number of sides, so this will be a regular polynomial.

Problem 18:

What is the generating function of the standard 6-sided die?

Problem 19:

What is the generating function of the die with sides $[1, 2, 3, 5]$?

Problem 20:

Let $A(x)$ and $B(x)$ be the generating functions of two dice.

What is the significance of $A(1)$?

Problem 21:

Using formulas we found earlier, show that the k^{th} coefficient of $A(x)B(x)$ is the number of ways to roll k as the sum of the two dice.

Problem 22:

Find a generating function for the sequence c_0, c_1, \dots , where c_k is the probability that the sum of the two dice is k .

Problem 23:

Using generating functions, find two six-sided dice whose sum has the same distribution as the sum of two standard six-sided dice.

That is, for any integer k , the number of ways that the sum of the two nonstandard dice rolls as k is equal to the number of ways the sum of two standard dice rolls as k . *Hint: factor polynomials.*

Part 4: Coins

Consider the following problem:

“How many different ways can you make change for \$0.50 using pennies, nickels, dimes, quarters and half-dollars?”

Most ways of solving this involve awkward brute-force approaches that don't reveal anything interesting about the problem: how can we change our answer if we want to make change for \$0.51, or \$1.05, or some other quantity?

We can use generating functions to solve this problem in a general way.

Definition 24:

Let p_0, p_1, p_2, \dots be such that p_k is the number of ways to make change for k cents with only pennies. Similarly, let...

- n_k be the number of ways to make change for k cents with only nickels;
- d_k be the number of ways using only dimes;
- q_k be the number of ways using only quarters;
- and h_k be the number of ways using only half-dollars.

Problem 25:

Let $p(x)$ be the generating function that corresponds to p_n . Express $p(x)$ as a rational function.

Problem 26:

Modify ?? to find expressions for $n(x)$, $d(x)$, $q(x)$, and $h(x)$.

Definition 27:

Now, let $N(x)$ be the generating function for the sequence n_0, n_1, \dots , where n_k is the number of ways to make change for k cents using pennies and nickels.

Similarly, let...

- let $D(x)$ be the generating function for the sequence using pennies, nickels, and dimes;
- let $Q(x)$ use pennies, nickels, dimes, and quarters;
- and let $H(x)$ use all coins.

Problem 28:

Express $N(x)$ as a rational function.

Problem 29:

Using the previous problem, write $D(x)$, then $Q(x)$, then $H(x)$ as rational functions.

Problem 30:

Using these generating functions, find recurrence relations for the sequences N_k, D_k, Q_k , and H_k .

Hint: Your recurrence relation for N_k should refer to the previous values of itself and some values of p_k . Your recurrence for D_k should refer to itself and N_k ; the one for Q_k should refer to itself D_k ; and the one for H_k should refer to itself and Q_k .

Problem 31:

Using these recurrence relations, fill following table and solve the original problem.

n	0	5	10	15	20	25	30	35	40	45	50
p_k											
N_k											
D_k											
Q_k											
H_k											

Part 5: Extra Problems

Problem 32: USAMO 1996 Problem 6

Determine (with proof) whether there is a subset X of the nonnegative integers with the following property: for any nonnegative integer n there is exactly one solution of $a + 2b = n$ with $a, b \in X$. (The original USAMO question asked about all integers, not just nonnegative - this is harder, but still approachable with generating functions.)

Problem 33: IMO Shortlist 1998

Let a_0, a_1, \dots be an increasing sequence of nonnegative integers such that every nonnegative integer can be expressed uniquely in the form $a_i + 2a_j + 4a_k$, where i, j, k are not necessarily distinct. Determine a_{1998} .

Problem 34: USAMO 1986 Problem 5

By a partition π of an integer $n \geq 1$, we mean here a representation of n as a sum of one or more positive integers where the summands must be put in nondecreasing order. (e.g., if $n = 4$, then the partitions π are $1 + 1 + 1 + 1$, $1 + 1 + 2$, $1 + 3$, $2 + 2$, and 4).

For any partition π , define $A(\pi)$ to be the number of ones which appear in π , and define $B(\pi)$ to be the number of distinct integers which appear in π (e.g., if $n = 13$ and π is the partition $1 + 1 + 2 + 2 + 2 + 5$, then $A(\pi) = 2$ and $B(\pi) = 3$).

Show that for any fixed n , the sum of $A(\pi)$ over all partitions of π of n is equal to the sum of $B(\pi)$ over all partitions of π of n .

Problem 35: USAMO 2017 Problem 2

Let m_1, m_2, \dots, m_n be a collection of n distinct positive integers. For any sequence of integers $A = (a_1, \dots, a_n)$ and any permutation $w = w_1, \dots, w_n$ of m_1, \dots, m_n , define an A -inversion of w to be a pair of entries w_i, w_j with $i < j$ for which one of the following conditions holds:

- $a_i \geq w_i > w_j$
- $w_j > a_i \geq w_i$
- $w_i > w_j > a_i$

Show that for any two sequences of integers $A = (a_1, \dots, a_n)$ and $B = (b_1, \dots, b_n)$ and for any positive integer k , the number of permutations of m_1, \dots, m_n having exactly k A -inversions is equal to the number of permutations of m_1, \dots, m_n having exactly k B -inversions. (The original USAMO problem allowed the numbers m_1, \dots, m_n to not necessarily be distinct.)